Center for Mathematical Physics BU-CMP/06-04

arXiv:math-ph/0611014v1 7 Nov 2006

FEYNMAN DIAGRAMS AND LAX PAIR EQUATIONS

GABRIEL BĂDIŢOIU AND STEVEN ROSENBERG

Abstract. We find a Lax pair equation corresponding to the Connes-Kreimer Birkhoff factorization of the character group of the Hopf algebra of Feynman diagrams. In particular, we obtain a flow for the character given by Feynman rules, and present a worked example.

1. Introduction

In the theory of integrable systems, a solution to a Lax pair equation associated to a coadjoint orbit of a semisimple Lie group is given by a Birkhoff factorization on the corresponding loop group. By the work of Connes-Kreimer [3], there is a Birkhoff factorization of characters on the Hopf algebra of Feynman diagrams. In this paper, we reverse the usual procedure in integrable systems by producing a Lax pair equation $\frac{dL}{dt} = [M, L]$ whose solution is given precisely by the Connes-Kreimer Birkhoff factorization (Theorem 5.6). The main technical issue, that the Lie algebra of infinitesimal characters is not semisimple, is overcome by passing to the double Lie algebra with the simplest possible Lie algebra structure. In particular, the Lax pair gives a flow for the character φ given by Feynman rules in dimensional regularization. It would be very interesting to know if this flow has physical significance.

In §§1-4, we introduce a method to produce a Lax pair on any Lie algebra from equations of motion on the double Lie algebra. In §5, we apply this method to the particular case of the Lie algebra of infinitesimal characters of the Hopf algebra of Feynman diagrams, and produce a Lax pair equation whose Birkhoff factorization coincides with the Connes-Kreimer factorization. In §6, we work out an explicit example of the theory on a finitely generated subalgebra of the Hopf algebra of Feynman diagrams.

It is natural to look for invariants of Lax pair equations by spectral curve techniques, and to linearize the flow on the Jacobian of the spectral curve. Unfortunately, in the worked example of §6, the spectral curve is highly reducible, and the only invariants we find are trivial. We hope to find examples with nontrivial invariants in the future.

2. The double Lie algebra and its associated Lie Group

There is a well known method to associate a Lax pair equation to a Casimir element on the dual $\mathfrak{g}^*$ of a semisimple Lie algebra $\mathfrak{g}$ [8]. The semisimplicity is used to produce an Ad-invariant, symmetric, non-degenerate bilinear form on $\mathfrak{g}$, allowing an identification

---

Date: February 7, 2008.

of $\mathfrak{g}$ with $\mathfrak{g}^*$. For a general Lie algebra $\mathfrak{g}$, there may be no Ad-invariant, symmetric, non-degenerate bilinear form on $\mathfrak{g}$. To produce a Lax pair, we need to extend $\mathfrak{g}$ to a larger Lie algebra with such a bilinear form. We do this by constructing a Lie bialgebra structure on $\mathfrak{g}$ and extending $\mathfrak{g}$ to $(\mathfrak{g}\oplus\mathfrak{g}^*, [\cdot,\cdot]_{\mathfrak{g}\oplus\mathfrak{g}^*})$, where $[\cdot,\cdot]_{\mathfrak{g}\oplus\mathfrak{g}^*}$ is the Lie bracket induced by the Lie bialgebra.

We recall the definition of a Lie bialgebra structure (see e.g. [6]).

Definition 2.1. A Lie bialgebra is a Lie algebra $(\mathfrak{g}, [\cdot,\cdot])$ with a linear map $\gamma : \mathfrak{g} \to \mathfrak{g}\otimes\mathfrak{g}$ such that

a) ${}^t\gamma : \mathfrak{g}^*\otimes\mathfrak{g}^* \to \mathfrak{g}^*$ defines a Lie bracket on $\mathfrak{g}^*$,

b) γ is a 1-cocycle of $\mathfrak{g}$, i.e.

$$\mathrm{ad}_x^{(2)}(\gamma(y)) - \mathrm{ad}_y^{(2)}(\gamma(x)) - \gamma([x,y]) = 0,$$

where $\mathrm{ad}_x^{(2)} : \mathfrak{g}^*\otimes\mathfrak{g}^* \to \mathfrak{g}^*\otimes\mathfrak{g}^*$ is given by $\mathrm{ad}_x^{(2)}(y\otimes z) = \mathrm{ad}_x(y)\otimes z + y\otimes\mathrm{ad}_x(z) = [x,y]\otimes z + y\otimes[x,z]$.

A Lie bialgebra $(\mathfrak{g}, [\cdot,\cdot], \gamma)$ induces an Lie algebra structure on the *double Lie algebra* $\mathfrak{g}\oplus\mathfrak{g}^*$ by

$$[X,Y]_{\mathfrak{g}\oplus\mathfrak{g}^*} = [X,Y],$$

$$[X^*,Y^*]_{\mathfrak{g}\oplus\mathfrak{g}^*} = {}^t\gamma(X\otimes Y),$$

$$[X,Y^*] = \mathrm{ad}^*_X(Y^*),$$

for X, $Y \in \mathfrak{g}$ and X^*, $Y^* \in \mathfrak{g}^*$, where ad^* is the coadjoint representation given by $\mathrm{ad}^*_X(Y^*)(Z) = -Y^*(\mathrm{ad}_X(Z))$ for $Z\in\mathfrak{g}$.

Since it is difficult to construct explicitly the Lie group associated to the Lie algebra $\mathfrak{g}\oplus\mathfrak{g}^*$, we will choose the trivial Lie bialgebra given by the cocyle $\gamma = 0$ and denote by $\delta = \mathfrak{g}\oplus\mathfrak{g}^*$ the associated Lie algebra. Let $\{Y_i, i = 1,\dots,l\}$ be a basis of $\mathfrak{g}$, with dual basis $\{Y_i^*\}$. The Lie bracket $[\cdot,\cdot]_\delta$ on δ is given by

$$[Y_i,Y_j]_\delta = [Y_i,Y_j],\ [Y_i^*,Y_j^*]_\delta = 0,\ [Y_i,Y_j^*]_\delta = -\sum_k c^j_{ik}Y_k^*,$$

where the c^j_{ik} are the structure constants: $[Y_i,Y_j] = \sum_k c^k_{ij}Y_k$.

The main point of this construction is that the natural pairing $\langle\cdot,\cdot\rangle : \delta\otimes\delta \to \mathbb{C}$ given by $\langle Y_i, Y_j^*\rangle = \delta_{ij}$ is an Ad-invariant symmetric non-degenerate bilinear form on the Lie algebra δ.

The Lie group naturally associated to δ is given by the following proposition.

Proposition 2.2. *Let* $\theta : g\times\mathfrak{g}^* \to \mathfrak{g}^*$ *be the coadjoint representation* $\theta(g,X) = \mathrm{Ad}^*_g(g)(X)$. *Then the Lie algebra of the semi-direct product* $\tilde{G} = g \rtimes_\theta \mathfrak{g}^*$ *is the double Lie algebra* δ.

Proof. The Lie group law on the semi-direct product $\tilde{G}$ is given by

$$(g,h)\cdot(g',h') = (gg', h+\theta_g(h')).$$

Let $\tilde{\mathfrak{g}}$ be the Lie algebra of $\tilde{G}$. Then the bracket on $\tilde{\mathfrak{g}}$ is given by

$$[X,Y^*]_{\tilde{\mathfrak{g}}} = d\theta(X,Y^*),\quad [X,Y]_{\tilde{\mathfrak{g}}} = [X,Y],\quad [X^*,Y^*]_{\tilde{\mathfrak{g}}} = 0,$$

for left-invariant vector fields X, Y of g and $X^*, Y^* \in \mathfrak{g}^*$. We have $d\theta(X,Y^*) = d\mathrm{Ad}_g(X,Y^*) = [X,Y^*]_\delta$ since $d\mathrm{Ad}_g = \mathrm{ad}_{\mathfrak{g}}$. $\square$

3. THE LOOP ALGEBRA OF A LIE ALGEBRA

Following [1], we consider the loop algebra

$$L\delta = \{L(\lambda) = \sum_{j=M}^{N} \lambda^j L_j \mid M,N\in\mathbb{Z}, L_j\in\delta\}.$$

The natural Lie bracket on $L\delta$ is given by

$$\left[\sum \lambda^i L_i, \sum \lambda^j L'_j\right] = \sum_k \lambda^k \sum_{i+j=k}[L_i, L'_j].$$

Set

$$\begin{aligned} L\delta_+ &= \{L(\lambda) = \sum_{j=0}^{N}\lambda^j L_j \mid N\in\mathbb{Z}^+\cup\{0\}, L_j\in\delta\}\\ L\delta_- &= \{L(\lambda) = \sum_{j=-M}^{-1}\lambda^j L_j \mid M\in\mathbb{Z}^+, L_j\in\delta\}.\end{aligned}$$

Let $P_+ : L\delta \to L\delta_+$ and $P_- : L\delta\to L\delta_-$ be the natural projections and set $R = P_+ - P_-$.

The natural pairing $\langle\cdot,\cdot\rangle$ on δ extends to an Ad-invariant, symmetric, non-degenerate pairing on $L\delta$ by setting

$$\left\langle \sum_{i=M}^{N}\lambda^i L_i, \sum_{j=M'}^{N'}\lambda^j L'_j\right\rangle = \sum_{i+j=-1}\langle L_i, L'_j\rangle.$$

For our choice of basis $\{Y_i\}$ of $\mathfrak{g}$, we get an isomorphism

$$I : L\delta^* \to L\delta \tag{3.1}$$

with

$$I\left(\sum L_i^j Y_j \lambda^i\right) = \sum L_i^j Y_j^* \lambda^{-1-i},$$

We will need the following lemmas.

Lemma 3.1. [1] *We have the following natural identifications:*

$$L\delta_+ = L\delta_-^* \text{ and } L\delta_- = L\delta_+^*.$$

Lemma 3.2. [8, Lem. 4.1] *Let φ be an* Ad*-invariant polynomial on δ. Then*

$$\varphi_{m,n}[L(\lambda)] = \mathrm{Res}_{\lambda=0}(\lambda^{-n}\varphi(\lambda^m L(\lambda))$$

is an Ad*-invariant polynomial on $L\delta$ for $m, n \in \mathbb{Z}$.*

As a double Lie algebra, δ has an Ad-invariant polynomial, the quadratic polynomial

$$\psi(Y) = \langle Y, Y\rangle$$

associated to the natural pairing. Let $Y_{l+i} = Y_i^*$ for $i \in \{1, \dots, l\}$, so elements of $L\delta$ can be written $L(\lambda) = \sum_{j=1}^{2l}\sum_{i=-M}^{N} L_i^j Y_j \lambda^i$. Then the Ad-invariant polynomials

$$\psi_{m,n}(L(\lambda)) = \mathrm{Res}_{\lambda=0}(\lambda^{-n}\psi(\lambda^m L(\lambda)), \tag{3.2}$$

defined as in Lemma 3.2 are given by

$$\psi_{m,n}(L(\lambda)) = 2\sum_{j=1}^{l}\sum_{i+k-n+2m=-1} L_i^j L_k^{j+l}. \tag{3.3}$$

Note that powers of ψ are also Ad-invariant polynomials on δ, so

$$\psi_{m,n}^k(L(\lambda)) = \mathrm{Res}_{\lambda=0}(\lambda^{-n}\psi^k(\lambda^m L(\lambda)) \tag{3.4}$$

are Ad-invariant polynomials on $L\delta$.

It would be interesting to classify all Ad-invariant polynomials on $L\delta$ in general.

4. The Lax pair equation

From [8, Theorem 2.1], the equation of motion induced by a Casimir function φ on the dual of a Lie algebra $\mathfrak{h}$ is given by

$$\frac{dL}{dt} = -\mathrm{ad}_{\mathfrak{h}}^* M \cdot L, \tag{4.1}$$

where $L \in \mathfrak{h}^*, M = \frac{1}{2}R(d\varphi(L)) \in \mathfrak{h}, R = P_+ - P_-$, where P_+, P_- are endomorphisms of $\mathfrak{h}$ such that

$$[X, Y]_R = [P_+X, P_+Y] - [P_-X, P_-Y]$$

is a Lie bracket on $\mathfrak{h}$.

Now we take $\mathfrak{h} = (L\delta)^* = L(\delta^*)$. Let $P_\pm$ be the projections of $L\delta^*$ onto $L\delta_\pm^*$. After identifying $L\delta^* = L\delta$ and $\mathrm{ad}^* = \mathrm{ad}$ via the map I in (3.1), the equation of motion (4.1) can be written in Lax pair form

$$\frac{dL}{dt} = [L, M], \tag{4.2}$$

where $M = \frac{1}{2}R(I(d\varphi(L(\lambda)))) \in L\delta$, and φ is a Casimir function on $L\delta^* = L\delta$.

Finding a solution for (4.2) reduces to the Riemann-Hilbert (or Birkhoff) factorization problem. The following theorem is a corollary of [8, Theorem 2.2].

Theorem 4.1. *Let φ be a Casimir function on $L\delta$ and set $X = I(d\varphi(L(\lambda))) \in L\delta$, for $L(\lambda) = L(0)(\lambda) \in L\delta$. Let $g_\pm(t)$ be the smooth curves in $L(\mathrm{Lie}(\delta))$ which solve the factorization problem*

$$\exp(-tX) = g_-(t)^{-1}g_+(t),$$

with $g_\pm(0) = e$, and with $g_+(t) = g_+(t)(\lambda)$ holomorphic in $\lambda \in \mathbb{C}$ and $g_-(t)$ a polynomial in $1/\lambda$ with no constant term. Let $M = \frac{1}{2}R(I(d\varphi(L(\lambda)))) \in L\delta$. Then the integral curve $L(t)$ of the Lax pair equation

$$\frac{dL}{dt} = [M, L]$$

is given by

$$L(t) = \mathrm{Ad}^*_{L\tilde{G}}g_\pm(t) \cdot L(0).$$

This Lax pair equation projects to a Lax pair equation on the loop algebra of the original Lie algebra $\mathfrak{g}$. Let π_1 be either the projection of $\tilde{G}$ onto G or its differential from δ onto $\mathfrak{g}$. This extends to a projection of $L\delta$ onto $L\mathfrak{g}$. The projection of (4.2) onto $L\mathfrak{g}$ is

$$\frac{d(\pi_1(L(t))}{dt} = [\pi_1(L), \pi_1(M)], \tag{4.3}$$

since $\pi_1 = d\pi_1$ commutes with the bracket. Thus the equations of motion (4.2) induce a Lax pair equation on $L\mathfrak{g}$, although this is not the equation of motion for a Casimir on $\mathfrak{g}^*$.

Theorem 4.2. *The Lax pair equation of Theorem 4.1 projects to a Lax pair equation on $L\mathfrak{g}$.*

When $\psi_{m,n}$ is the Casimir function on $L\delta$ given by (3.2), X can be written nicely in terms of $L(\delta)$.

Proposition 4.3. *Let $X = I(d\psi_{m,n}(L(\lambda)))$. Then*

$$X = 2\lambda^{-n+2m}L(\lambda). \tag{4.4}$$

Proof. Write $L(\lambda) = \sum_{i,j} L_i^j \lambda^i Y_j$. By formula (3.3), we have

$$\frac{\partial \psi_{m,n}}{\partial L_p^t} = \begin{cases} 2L_{n-1-2m-p}^{t+l}, & \text{if } t \leq l \\ 2L_{n-1-2m-p}^{t-l}, & \text{if } t > l. \end{cases} \tag{4.5}$$

Therefore

$$\begin{aligned} X &= I(d\psi_{m,n}(L(\lambda))) = \sum_{p,t} \frac{\partial \psi_{m,n}}{\partial L_p^t}\lambda^{-1-p}Y_t^* \\ &= 2\lambda^{-n+2m}\sum_p(\sum_{t=1}^{l} L_{n-1-2m-p}^{t+l}Y_{t+l}\lambda^{n-1-2m-p} + \sum_{t=l+1}^{2l} L_{n-1-2m-p}^{t-l}Y_{t-l}\lambda^{n-1-2m-p}) \\ &= 2\lambda^{-n+2m}L(\lambda). \end{aligned}$$

□

5. The main theorem for Hopf algebras

Let $H = (H, 1, \mu, \Delta, \varepsilon, S)$ be a Hopf algebra over $\mathbb{C}$ and let $H^* = (H^*, 1_*, \mu_* = \Delta^t, \Delta_* = \mu^t, \varepsilon_* = 1^t, S^* = S^t)$ be the algebraic dual of H.

Definition 5.1. The **character group** $\mathrm{Char}(H)$ of a Hopf algebra H is the group of "group-like elements" of H^* (see [5]):

$$G = \{\phi \in H^* \mid \phi \neq 0,\ \Delta_*\phi = \phi \otimes \phi\}.$$

The group law is given by the convolution product and the unit element is 1_*:

$$(\psi_1 \star \psi_2)(h) = \langle \psi_1 \otimes \psi_2, \Delta h \rangle,$$

$$\langle 1_*, h \rangle = \varepsilon(h), \quad \text{for } h \in H.$$

Definition 5.2. An **infinitesimal character** of a Hopf algebra H is a $\mathbb{C}$-linear map $Z : H \to \mathbb{C}$ satisfying

$$\langle Z, hk \rangle = \langle Z, h \rangle \varepsilon(k) + \varepsilon(h) \langle Z, k \rangle.$$

The set of infinitesimal characters is denoted by $\partial\mathrm{Char}(H)$ and is endowed with a Lie algebra bracket:

$$[Z, Z'] = Z \star Z' - Z' \star Z, \quad \text{for } Z,\ Z' \in \partial Char(H),$$

where $\langle Z \star Z', h \rangle = \langle Z \otimes Z', \Delta(h) \rangle$.

Let $(H, 1 = \emptyset, \mu, \Delta, \varepsilon, S)$ be the universal Hopf algebra of all Feynman graphs. For computations later, we also consider a Hopf subalgebra H_0 generated by a finite number of Feynman graphs $A_1, A_2, \ldots, A_{l-1}, A_l = \emptyset$, and let H_1 be the Hopf subalgebra generated by $A_1, \ldots, A_{l-1}$. Let G be the Lie group of characters of H, and let G_0 be the Lie group of characters of H_0. The Lie algebra of infinitesimal characters $\mathfrak{g}, \mathfrak{g}_0$ of H, H_0 are precisely the Lie algebras of G, G_0, respectively.

Let Z_0 be the infinitesimal character of H given by $Z_0(h) = \varepsilon(h)$. For any generator of H, viz. $T \in \{A_1, \ldots A_{l-1}, A_l = \emptyset\}$, let Z_T be the infinitesimal character given by $Z_T(T') = \delta_{T,T'}$. Notice that $Z_0 = Z_\emptyset$. The Lie algebra $\mathfrak{g}_0$ is generated by $Z_0,\ Z_{A_1}, \ldots,\ Z_{A_{l-1}}$. Let $\mathfrak{g}_1$ be the Lie subalgebra of $\mathfrak{g}_0$ generated by $Z_{A_1},\ Z_{A_2}, \ldots Z_{A_{l-1}}$. Let G_1 be the Lie subgroup of G_0 corresponding to $\mathfrak{g}_1$. Set $Y_i = Z_{A_i}$ for $i \in \{1, \ldots, l-1\}$ and set $Y_l = Z_0$. We also let $\mathfrak{g}_1$ denote the Lie subalgebra of $\mathfrak{g}$ with basis $Z_T, T \neq \emptyset$, and let G_1 also denote the corresponding subgroup of G.

Remark 5.3. We have $\mathfrak{g} = \mathfrak{g}_1 \oplus \langle Z_\emptyset \rangle$. At the Lie group level, G_0 is the semi-direct product $G_1 \rtimes \mathbb{C}$ given by

$$(g, t) \cdot (g', t') = (g \cdot \theta_t(g'), t + t'),$$

where $\theta_t(g)(\Gamma) = e^{t\#(\Gamma)}g(\Gamma)$ for $\Gamma \in H_1$, and $\#(\Gamma)$ is the number of independent loops of Γ.

Recall that for any Lie group K, an element $L(\lambda) \in LK$ has a Birkhoff decomposition if $L(\lambda) = L(\lambda)_-^{-1}L(\lambda)_+$ with $L(\lambda)_-^{-1}$ holomorphic in $\lambda^{-1} \in \mathbb{P} - \{0\}$ and $L(\lambda)_+$ holomorphic in $\lambda \in \mathbb{P} - \{\infty\}$. In the next lemma, $\tilde{G}$ refers either to $G_0 \rtimes_\theta \mathfrak{g}_0^*$ as in Prop. 2.2 or to $G \rtimes_\theta \mathfrak{g}^*$.

Lemma 5.4. *Let* (g, α) *be an element in* $L\tilde{G}$. *If* $(g, \alpha) = (g_-, \alpha_-)^{-1}(g_+, \alpha_+)$ *then* $g = g_-^{-1}g_+$ *and* $\alpha = \mathrm{Ad}^*(g_-^{-1})(-\alpha_- + \alpha_+)$.

Proof. We recall that $(g_1, \alpha_1)(g_2, \alpha_2) = (g_1g_2, \alpha_1 + \mathrm{Ad}^*(g_1)(\alpha_2))$. Notice that $(g_-, \alpha_-)^{-1} = (g_-^{-1}, -\mathrm{Ad}^*(g_-^{-1})(\alpha_-))$, so $(g_-, \alpha_-)^{-1}(g_+, \alpha_+) = (g_-^{-1}g_+, -\mathrm{Ad}^*(g_-^{-1})(\alpha_-) + \mathrm{Ad}^*(g_-^{-1})(\alpha_+)$. □

We prove the existence of a Birkhoff decomposition for any element $(g, \alpha) \in L\tilde{G}$.

Theorem 5.5. *Every* $(g, \alpha) \in L\tilde{G}$ *has a Birkhoff decomposition* $(g, \alpha) = (g_-, \alpha_-)^{-1}(g_+, \alpha_+)$ *with* (g_+, α_+) *a polynomial in* λ *and* (g_-, α_-) *a polynomial in* λ^{-1} *without constant term.*

Proof. Let $g = g_-^{-1}g_+$ be the Birkhoff decomposition of g in LG_0 given in [4]. Let $\alpha_+ = P_+(\mathrm{Ad}^*(g_-)(\alpha))$ and $\alpha_- = -P_-(\mathrm{Ad}^*(g_-)(\alpha))$. Then, by Lemma 5.4, $(g, \alpha) = (g_-, \alpha_-)^{-1}(g_+, \alpha_+)$. □

In [3], Connes and Kreimer give a Birkhoff decomposition for the character group of the Feynman graph Hopf algebra, and in particular for the normalized loop character $\bar{\varphi}(\lambda, q)$ of dimensional regularization. Here

$$\bar{\varphi}(\lambda, q) = \frac{\varphi(\lambda)}{q^2},$$

where $\varphi(\lambda, q)$ is the usual character given by dimensional regularization and Feynman rules. We consider the algebra of formal power series

$$\Omega\delta = \{L(\lambda) = \sum_{j=-\infty}^{\infty} \lambda^j L_j \mid L_j \in \delta\}.$$

The natural Lie bracket on $\Omega\delta$ is

$$\left[\sum \lambda^i L_i, \sum \lambda^j L'_j\right] = \sum_k \lambda^k \sum_{i+j=k} [L_i, L'_j].$$

Set

$$\begin{aligned} \Omega\delta_+ &= \{L(\lambda) = \sum_{j=0}^{\infty} \lambda^j L_j \mid L_j \in \delta\} \\ \Omega\delta_- &= \{L(\lambda) = \sum_{j=-\infty}^{-1} \lambda^j L_j \mid L_j \in \delta\}. \end{aligned}$$

Recall that π_1 denotes either the projection of the double Lie group $\tilde{G}$ to its first factor G, its differential, or its extention to the loop group and loop algebra. We denote the image of an element by adding a tilde, e.g. $\pi_1(L(\lambda)) = \tilde{L}(\lambda)$.

Theorem 5.6. *Let H be the Hopf algebra of Feynman graphs, and let $\bar{\varphi}(\lambda) \in \mathrm{Char}(H) = G$. Set*

$$L_0(\lambda) = \frac{1}{2}\lambda^{n-2m}\exp^{-1}(\bar{\varphi}(\lambda)) \quad \textit{and} \quad X = I(d\psi_{m,n}(L_0(\lambda)),$$

where $\psi_{m,n}$ is the Casimir function on $\Omega\delta$ given by

$$\psi_{m,n}(L(\lambda)) = \mathrm{Res}_{\lambda=0}(\lambda^m\langle\lambda^n L(\lambda), \lambda^n L(\lambda)\rangle).$$

Then $\exp(X) = \bar{\varphi}$ and the solution of

$$\frac{dL}{dt} = [M, L], \quad M = \frac{1}{2}R(I(d\psi_{m,n}(L(\lambda))) \tag{5.1}$$

with the initial condition $L(0) = L_0$ is given by

$$L(t) = \mathrm{Ad}^*_{L\tilde{G}}g_\pm(t)\cdot L_0, \tag{5.2}$$

with $\exp(-tX) = g_-(t)^{-1}g_+(t)$. Applying π_1, we get a Lax pair equation

$$\frac{d\tilde{L}}{dt} = [\tilde{M}, \tilde{L}]$$

on LG, with solution $\tilde{L}(t) = \mathrm{Ad}^*_{LG}\tilde{g}_\pm(t)\cdot\tilde{L}_0$. In particular,

$$\bar{\varphi} = \exp(X) = \tilde{g}_-(-1)^{-1}\tilde{g}_+(-1)$$

is the Connes-Kreimer factorization of $\bar{\varphi}$. The same results hold for any finitely generated Hopf subalgebra H_0 of H.

Proof. By [7, p. 32], the exponential map $\exp : \mathfrak{g} \to G$ is bijective, so $L_0(\lambda)$ exists. The theorem then follows from Theorem 4.1 and (4.3) applied to the natural pairing on δ as in (3.2) and the uniqueness of the Birkhoff factorization. □

Remark 5.7. We have to be careful passing from finite dimensional Lie algebras to the infinite dimensional Lie algebra $\mathfrak{g}$ of infinitesimal characters on H, since the finite dimensional identification of a Lie algebra $\mathfrak{h}$ with $\mathfrak{h}^*$ via a choice of basis of $\mathfrak{h}$ may not be valid. However, we just need the part of $\mathfrak{g}^*$ spanned by the dual basis to a fixed basis of $\mathfrak{g}$, so the finite dimensional theory extends.

6. A WORKED EXAMPLE

In this section we discuss a specific example of the main theorem.

Let

$A =$  , $AA =$  , $AAA =$  , $AAAA =$  , $B =$  .

Let H_0 be the Hopf subalgebra generated by $\emptyset$, A, AA, AAA , $AAAA$, B, and let H_1 be the Hopf subalgebra generated by A, AA, AAA , $AAAA$, B. Let G_0, G_1 be the Lie groups of characters of H_0, H_1, respectively. The Lie algebra of infinitesimal characters $\mathfrak{g}_0$ of H_0 is the Lie algebra of G_0, and similarly for $\mathfrak{g}_1$. In particular, the Lie algebra $\mathfrak{g}_1$ of G_1 is generated by $Z_A,\ Z_{AA},\ Z_{AAA},\ Z_{AAAA},\ Z_B$. We identify G_1 with $\mathbb{C}^5$ using the normal coordinates defined in [2]. The group law induced on $\mathbb{C}^5$ is given by the following lemma.

Lemma 6.1. *Let* $\oplus : \mathbb{C}^5 \times \mathbb{C}^5 \to \mathbb{C}^5$ *be the group law on* $\mathbb{C}^5$ *given by*

$$(x_1, x_2, x_3, x_4, x_5) \oplus (y_1, y_2, y_3, y_4, y_5) = (x_1+y_1, x_2+y_2, x_3+y_3, x_4+y_4+x_1y_2-x_2y_1, x_5+y_5).$$

Define $F : G_1 \to (\mathbb{C}^5, \oplus)$ *by*

$$F(\varphi) = (\varphi(A), \varphi(AA) - \frac{1}{2}\varphi(A)^2, \varphi(AAA) - \varphi(A)\varphi(AA) + \frac{1}{6}\varphi(A)^3,$$

$$\varphi(B)-\varphi(A)\varphi(AA)+\frac{1}{6}\varphi(A)^3, \varphi(AAAA)-\varphi(A)\varphi(AAA)-\frac{1}{2}\varphi(AA)^2+\varphi(A)^2\varphi(AA)-\frac{1}{4}\varphi(A)^4).$$

Then F *is a group isomorphism.*

Proof. Let $(F_1, F_2, F_3, F_4, F_5) = F$. We have

$$\Delta(A) = A \otimes 1 + 1 \otimes A,$$

$$\Delta(AA) = AA \otimes 1 + 1 \otimes AA + A \otimes A,$$

$$\Delta(AAA) = AAA \otimes 1 + 1 \otimes AAA + AA \otimes A + A \otimes AA,$$

$$\Delta(AAAA) = AAAA \otimes 1 + 1 \otimes AAAA + AAA \otimes A + A \otimes AAA + AA \otimes AA,$$

$$\Delta(B) = B \otimes 1 + 1 \otimes B + 2A \otimes AA.$$

Therefore

$$(\phi_1 \star \phi_2)(A) = \langle \phi_1 \otimes \phi_2, \Delta(A) \rangle = \phi_1(A) + \phi_2(A),$$

$$\begin{aligned} (\phi_1 \star \phi_2)(AA) - \frac{1}{2}(\phi_1 \star \phi_2)(A)^2 &= \langle \phi_1 \otimes \phi_2, \Delta(AA) \rangle - \frac{1}{2}(\phi_1 \star \phi_2)(A)^2 \\ &= \phi_1(AA) + \phi_2(AA) + \phi_1(A)\phi_2(A) - \frac{1}{2}(\phi_1(A) + \phi_2(A))^2 \\ &= (\phi_1(AA) - \frac{1}{2}\phi_1(A)^2) + (\phi_2(AA) - \frac{1}{2}\phi_2(A)^2), \end{aligned}$$

which implies $F_k(\phi_1 \star \phi_2) = F_k(\phi_1) + F_k(\phi_2)$ for $k \in \{1, 2\}$. By direct computations we also get

$$F_k(\phi_1 \star \phi_2) = F_k(\phi_1) + F_k(\phi_2) \ \text{ for } k \in \{3, 5\},$$

$$F_4(\phi_1 \star \phi_2) = F_4(\phi_1) + F_4(\phi_2) + F_1(\phi_1)F_2(\phi_2) - F_2(\phi_1)F_1(\phi_2).$$

□

Using this identification of G_1 with $(\mathbb{C}^5, \oplus)$, we can identify G_0 with $(\mathbb{C}^6, \oplus)$, where

$$(x_1, x_2, x_3, x_4, x_5, t) \oplus (y_1, y_2, y_3, y_4, y_5, t') =$$
$$(x_1 + e^t y_1, x_2 + e^{2t} y_2, x_3 + e^{3t} y_3, x_4 + e^{3t} y_4 + e^{2t} x_1 y_2 - e^t x_2 y_1, x_5 + e^{4t} y_5, t + t').$$

The following lemma gives a basis of the left invariant vector fields on G_0 and the structure constants of $\mathfrak{g}_0$.

Lemma 6.2. *Let*

$$Y_1 = e^t \left(\frac{\partial}{\partial y_1} - y_2 \frac{\partial}{\partial y_4} \right), \qquad Y_2 = e^{2t} \left(\frac{\partial}{\partial y_2} + y_1 \frac{\partial}{\partial y_4} \right),$$
$$Y_3 = e^{3t} \frac{\partial}{\partial y_3}, \qquad Y_4 = e^{3t} \frac{\partial}{\partial y_4}, \qquad Y_5 = e^{4t} \frac{\partial}{\partial y_5}, \qquad Z_0 = \frac{\partial}{\partial t}.$$

where $(y_1, y_2, y_3, y_4, y_5, t)$ *are the standard coordinates on* $\mathbb{C}^5$. *Then* $\{Y_1, Y_2, Y_3, Y_4, Y_5, Z_0\}$ *is a basis of the left invariant vector fields on* G_0. *We have* $[Y_i, Y_j] = 0$ *for any* $(i, j) \neq (1, 2), (2, 1)$, *and* $[Y_1, Y_2] = -[Y_2, Y_1] = 2Y_4$, $[Z_0, Y_1] = Y_1$, $[Z_0, Y_2] = 2Y_2$, $[Z_0, Y_3] = 3Y_3$, $[Z_0, Y_4] = 3Y_4$ *and* $[Z_0, Y_5] = 4Y_5$.

Proof. This follows from the easily computed formulas

$$L_x \left(\frac{\partial}{\partial y_1} \right) = e^t \left(\frac{\partial}{\partial y_1} - x_2 \frac{\partial}{\partial y_4} \right), \quad L_x \left(\frac{\partial}{\partial y_2} \right) = e^{2t} \left(\frac{\partial}{\partial y_2} + x_1 \frac{\partial}{\partial y_4} \right),$$
$$L_x \left(\frac{\partial}{\partial y_3} \right) = e^{3t} \left(\frac{\partial}{\partial y_3} \right), \quad L_x \left(\frac{\partial}{\partial y_4} \right) = e^{3t} \left(\frac{\partial}{\partial y_4} \right), \quad L_x \left(\frac{\partial}{\partial y_5} \right) = e^{4t} \left(\frac{\partial}{\partial y_5} \right),$$
$$L_x \left(\frac{\partial}{\partial t'} \right) = \frac{\partial}{\partial t'}.$$

□

6.1. **The exponential map and the adjoint and coadjoint representations.**

Lemma 6.3. *The exponential* exp : $\mathfrak{g}_0 \to G_0$ *is given by*

$$\exp(a_1 Y_1 + a_2 Y_2 + a_3 Y_3 + a_4 Y_4 + a_5 Y_5 + a_6 Z_0) =$$
$$= \begin{cases} (a_1, a_2, a_3, a_4, a_5, 0), & \textit{if } a_6 = 0 \\ \left(\frac{a_1(e^{a_6}-1)}{a_6}, \frac{a_2(e^{2a_6}-1)}{2a_6}, \frac{a_2(e^{3a_6}-1)}{3a_6}, \frac{a_4(e^{3a_6}-1)}{3a_6} + \frac{a_1 a_2}{2a_6^2}(\frac{e^{3a_6}-1}{3} - e^{2a_6 t} + e^{a_6 t}), \frac{a_5(e^{4a_6}-1)}{4a_6}, a_6 \right), & \textit{if } a_6 \neq 0, \end{cases}$$

and exp *is bijective.*

Proof. Let $Y \in \mathfrak{g}_0$ and let $\gamma(t)$ be the 1-parameter subgroup of G_0 generated by Y. Set

$$\begin{aligned} Y &= a_1 Y_1 + a_2 Y_2 + a_3 Y_3 + a_4 Y_4 + a_5 Y_5 + a_6 Z_0, \\ \gamma(t) &= (g_1(t), g_2(t), g_3(t), g_4(t), g_5(t), g_6(t)). \end{aligned}$$

To find $\gamma(t) = \exp(tY)$, we solve the differential equation

$$L_{\gamma(t)^{-1}} \dot{\gamma}(t) = a_1 Y_1 + a_2 Y_2 + a_3 Y_3 + a_4 Y_4 + a_5 Y_5 + a_6 Z_0$$

with the initial condition $\gamma(0) = 0$. First notice that

$$\begin{aligned} L_{\gamma(t)^{-1}}\dot{\gamma}(t) &= \dot{g}_1 e^{-g_6}\frac{\partial}{\partial h_1} + \dot{g}_2 e^{-2g_6}\frac{\partial}{\partial h_2} + \dot{g}_3 e^{-3g_6}\frac{\partial}{\partial h_3} + e^{-3g_6}(\dot{g}_1 g_2 - \dot{g}_2 g_1 + \dot{g}_4)\frac{\partial}{\partial h_4} \\ &\quad + \dot{g}_5 e^{-4g_6}\frac{\partial}{\partial h_5} + \dot{g}_6\frac{\partial}{\partial h_6}. \end{aligned}$$

Then $a_k = \dot{g}_k e^{-kg_6}$ for $k \in \{1,2,3\}$, $a_5 = \dot{g}_5 e^{-4g_6}$, $a_6 = \dot{g}_6$ and $a_4 = (\dot{g}_4 + \dot{g}_1 g_2 - \dot{g}_2 g_1)e^{-3g_6}$, with the initial conditions $g_1(0) = g_2(0) = g_3(0) = g_4(0) = g_5(0) = g_6(0) = 0$. Therefore $g_6 = a_6 t$, $g_k(t) = a_k(e^{ka_6 t} - 1)/(ka_6)$ for $k \in \{1,2,3\}$, $g_5(t) = a_5(e^{4a_6 t} - 1)/(4a_6)$ and

$$g_4 = \frac{a_4(e^{3a_6 t} - 1)}{3a_6} - \int \frac{a_1 e^{a_6 t} a_2(e^{2a_6 t} - 1)}{2a_6}\, dt + \int \frac{a_2 e^{2a_6 t} a_1(e^{a_6 t} - 1)}{a_6}\, dt.$$

$$g_4(t) = \frac{a_4(e^{3a_6 t} - 1)}{3a_6} + \frac{a_1 a_2}{2a_6^2}\left(\frac{e^{3a_6 t} - 1}{3} - e^{2a_6 t} + e^{a_6 t}\right)$$

If $a_6 \neq 0$ then

$$\begin{aligned} \exp\left(\sum_{k=1}^{5} a_k Y_k + a_6 Z_0\right) = \Bigg(&\frac{a_1(e^{a_6} - 1)}{a_6}, \frac{a_2(e^{2a_6} - 1)}{2a_6}, \frac{a_2(e^{3a_6} - 1)}{3a_6}, \\ &\frac{a_4(e^{3a_6} - 1)}{3a_6} + \frac{a_1 a_2}{2a_6^2}\left(\frac{e^{3a_6} - 1}{3} - e^{2a_6 t} + e^{a_6 t}\right), \frac{a_5(e^{4a_6} - 1)}{4a_6}, a_6 \Bigg). \end{aligned}$$

If $a_6 = 0$ then

$$\exp\left(\sum_{k=1}^{5} a_k Y_k\right) = (a_1, a_2, a_3, a_4, a_5, 0).$$

□

The adjoint and coadjoint representations of G_0 are given by the following lemmas.

Lemma 6.4. *i) The adjoint representation* $\mathrm{Ad}_{G_0} : G_0 \to \mathrm{GL}(\mathfrak{g}_0)$ *is given by*

$$\mathrm{Ad}_{G_0}(g_1, \ldots, g_5, g_6) = \begin{pmatrix} e^{g_6} & 0 & 0 & 0 & 0 & -g_1 \\ 0 & e^{2g_6} & 0 & 0 & 0 & -2g_2 \\ 0 & 0 & e^{3g_6} & 0 & 0 & -3g_3 \\ -2g_2 e^{g_6} & 2g_1 e^{2g_6} & 0 & e^{3g_6} & 0 & -g_1 g_2 - 3g_4 \\ 0 & 0 & 0 & 0 & e^{4g_6} & -4g_5 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

ii) The coadjoint representation $\mathrm{Ad}^*_{G_0} : G_0 \to \mathrm{GL}(\mathfrak{g}^*_0)$ *is given by*

$$\mathrm{Ad}^*_{G_0}(g_1, \ldots, g_5, g_6) = \begin{pmatrix} e^{-g_6} & 0 & 0 & 2g_2 e^{-3g_6} & 0 & 0 \\ 0 & e^{-2g_6} & 0 & -2g_1 e^{-3g_6} & 0 & 0 \\ 0 & 0 & e^{-3g_6} & 0 & 0 & 0 \\ 0 & 0 & 0 & e^{-3g_6} & 0 & 0 \\ 0 & 0 & 0 & 0 & e^{-4g_6} & 0 \\ e^{-g_6} g_1 & 2e^{-2g_6} g_2 & 3e^{-3_6} g_3 & -e^{-3_6} g_1 g_2 + 3e^{-3g_6} g_4 & 4e^{-4g_6} g_5 & 1 \end{pmatrix}$$

Proof. To show i), we straightforwardly compute $\mathrm{Ad}_{G_0}(g) = dC_g$ where $C_g(h) = ghg^{-1}$, noting that

$$g^{-1} = (x_1, x_2, x_3, x_4, x_5, t)^{-1} = (-e^{-t}x_1, -e^{-2t}x_2, -e^{-3t}x_3, -e^{-3t}x_4, -e^{-4t}x_5, -t).$$

ii) then follows from *i)* and $\mathrm{Ad}^*(g) = (\mathrm{Ad}(g^{-1}))^t$. □

Lemma 6.5. *1) For the basis* $\{Y_1, \dots, Y_6\}$ *of Lemma 6.2,* $\mathrm{ad} : \mathfrak{g}_0 \to \mathfrak{gl}(\mathfrak{g}_0)$ *is given by*

$$\mathrm{ad}\left(\sum_{i=1}^{6} c_i Y_i\right) = \begin{pmatrix} c_6 & 0 & 0 & 0 & 0 & -c_1 \\ 0 & 2c_6 & 0 & 0 & 0 & -2c_2 \\ 0 & 0 & 3c_6 & 0 & 0 & -3c_3 \\ -2c_2 & 2c_1 & 0 & 3c_6 & 0 & -3c_4 \\ 0 & 0 & 0 & 0 & 4c_6 & -4c_5 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

2) $\mathrm{ad}^* : \mathfrak{g}_0 \to \mathfrak{gl}(\mathfrak{g}_0^*)$ *is given by*

$$\mathrm{ad}^*\left(\sum_{i=1}^{6} c_i Y_i\right) = \begin{pmatrix} -c_6 & 0 & 0 & 2c_2 & 0 & 0 \\ 0 & -2c_6 & 0 & -2c_1 & 0 & 0 \\ 0 & 0 & -3c_6 & 0 & 0 & 0 \\ 0 & 0 & 0 & -3c_6 & 0 & 0 \\ 0 & 0 & 0 & 0 & -4c_6 & 0 \\ c_1 & 2c_2 & 3c_3 & 3c_4 & 4c_5 & 0 \end{pmatrix}$$

Since $d\mathrm{Ad} = \mathrm{ad}$, the proof follows by differentiating the formulas in Lemma 6.4.

6.2. The double Lie algebra and its associated Lie group.

The conditions a) and b) in Definition 2.1 of a Lie bialgebra can be written in a basis as a system of quadratic equations. Solving this system explicitly, in our case via Mathematica, gives the following proposition.

Proposition 6.6. *There are* 43 *families of bialgebra structures* γ *on* $\mathfrak{g}$.

In fact, the system of quadratic equations involves 90 variables. Mathematica gives 1 solution with 82 linear relations (and so 8 degrees of freedom), 7 solutions with 83 linear relations, 16 solutions with 84 linear relations, 13 solutions with 85 linear relations, 5 solutions with 86 linear relations, and 1 solution with 87 linear relations.

As it is difficult to construct explicitly a Lie group corresponding to the Lie algebra $\mathfrak{g} \oplus \mathfrak{g}^*$ for an arbitrary choice of γ, we take the simplest choice $\gamma = 0$ and let $\tilde{G} = G_0 \rtimes \mathfrak{g}_0^*$ be the corresponding Lie group of $\delta = \mathfrak{g} \oplus \mathfrak{g}^*$.

Remark 6.7. The group law on $\tilde{G}$ is given by

$$((g_1, g_2, g_3, g_4, g_5, g_6), (h_1, h_2, h_3, h_4, h_5, h_6)) \cdot ((g_1', g_2', g_3', g_4', g_5', g_6'), (h_1', h_2', h_3', h_4', h_5', h_6')) =$$

$$(g_1 + e^{g_6} g_1', g_2 + e^{2g_6} g_2', g_3 + e^{3g_6} g_3', g_4 + e^{3g_6} g_4' + g^{2g_6} g_1 g_2' - g^{g_6} g_2 g_1', g_5 + e^{4g_6} g_5', g_6 + g_6',$$

$$h_1 + e^{g_6}h_1' - 2g_2e^{g_6}h_4', h_2 + e^{2g_6}h_2' + 2g_2e^{2g_6}h_4', h_3 + e^{3g_6}h_3', h_4 + e^{3g_6}h_4', h_5 + e^{4g_6}h_5',$$
$$h_6 - g_1h_1' - 2g_2h_2' - 3g_3h_3' - 3g_4h_4' - 4g_5h_5' + h_6').$$

6.3. **The adjoint representations** ad_δ **and** $\mathrm{Ad}_{\tilde{G}}$**.** Let $\mathrm{ad}_\delta : \delta \to \mathfrak{gl}(\delta)$ be the adjoint representation of δ. Computing ad_δ explicitly, for example via Mathematica, we get

$$\mathrm{ad}_\delta\left(\sum_{i=1}^{12} x_iY_i\right) =$$

$$\begin{pmatrix}
x_6 & 0 & 0 & 0 & 0 & -x_1 & 0 & 0 & 0 & 0 & 0 & 0\\
0 & 2x_6 & 0 & 0 & 0 & -2x_2 & 0 & 0 & 0 & 0 & 0 & 0\\
0 & 0 & 3x_6 & 0 & 0 & -3x_3 & 0 & 0 & 0 & 0 & 0 & 0\\
-2x_2 & 2x_1 & 0 & 3x_6 & 0 & -3x_4 & 0 & 0 & 0 & 0 & 0 & 0\\
0 & 0 & 0 & 0 & 4x_6 & -4x_5 & 0 & 0 & 0 & 0 & 0 & 0\\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0\\
0 & -2x_{10} & 0 & 0 & 0 & x_7 & -x_6 & 0 & 0 & 2x_2 & 0 & 0\\
2x_{10} & 0 & 0 & 0 & 0 & 2x_8 & 0 & -2x_6 & 0 & -2x_1 & 0 & 0\\
0 & 0 & 0 & 0 & 0 & 3x_9 & 0 & 0 & -3x_6 & 0 & 0 & 0\\
0 & 0 & 0 & 0 & 0 & 3x_{10} & 0 & 0 & 0 & -3x_6 & 0 & 0\\
0 & 0 & 0 & 0 & 0 & 4x_{11} & 0 & 0 & 0 & 0 & -4x_6 & 0\\
-x_7 & -2x_8 & -3x_9 & -3x_{10} & -4x_{11} & 0 & x_1 & 2x_2 & 3x_3 & 3x_4 & 4x_5 & 0
\end{pmatrix}$$

where $Y_{6+t} = Y_t$ for $t \in \{1, \ldots, 6\}$.

Corollary 6.8. $\mathrm{Ker}(\mathrm{ad}_\delta) = \mathrm{Span}\{Y_{12}\}$.

The adjoint and coadjoint representations of $\tilde{G}$ are given in the following proposition.

Proposition 6.9.

$$\mathrm{Ad}_{\tilde{G}}(g_1, g_2, \ldots, g_{12}) = \begin{pmatrix} \mathrm{Ad}_{G_0}(g_1, g_2, \ldots, g_6) & 0 \\ H(g_1, g_2, \ldots, g_{12}) & \mathrm{Ad}^*_{G_0}(g_1, g_2, \ldots, g_6) \end{pmatrix}$$

$$\mathrm{Ad}^*_{\tilde{G}}(g_1, g_2, \ldots, g_{12}) = \begin{pmatrix} \mathrm{Ad}^*_{G_0}(g_1, g_2, \ldots, g_6) & H(g_1, g_2, \ldots, g_{12}) \\ 0 & \mathrm{Ad}_{G_0}(g_1, g_2, \ldots, g_6) \end{pmatrix}$$

where $H(g_1, g_2, \ldots, g_{12})$ *is a* 6×6 *matrix given by*

$$\begin{pmatrix}
0 & -2e^{2g_6}g_{10} & 0 & 0 & 0 & 4g_{10}g_2 + g_7\\
2e^{g_6}g_{10} & 0 & 0 & 0 & 0 & -2g_1g_{10} + 2g_8\\
0 & 0 & 0 & 0 & 0 & 3g_9\\
0 & 0 & 0 & 0 & 0 & 3g_{10}\\
0 & 0 & 0 & 0 & 0 & 4g_{11}\\
6e^{g_6}g_{10}g_2 - e^{g_6}g_7 & -6e^{2g_6}g_1g_{10} - 2e^{2g_6}g_8 & -3e^{3g_6}g_9 & -3e^{3g_6}g_{10} & -4e^{4g_6}g_{11} & z
\end{pmatrix}$$

and

$$z = 3g_1g_{10}g_2 + 9g_{10}g_4 + 16g_{11}g_5 + g_1g_7 + 4g_2g_8 + 9g_3g_9.$$

The proof is straightforward.

6.4. **Some $\mathrm{Ad}_{\tilde{G}}$-invariant polynomials on δ and $\mathbf{L}(\delta)$.** We note that $\mathrm{Tr}(\mathrm{ad}(a)^k), k \in \mathbb{Z}^+$, are Ad-invariant polynomials on δ. By Lemma 3.2, these induce Ad-invariant polynomials on $L\delta$, i.e. Casimir functions (constants of motions) on $L\delta$. Explicit computations give the following lemma.

Lemma 6.10. *Let $\varphi : \delta \to \mathbb{C}$ be the map given by $\varphi(a) = \mathrm{Tr}(\mathrm{ad}(a)^k)$. For odd positive integers k, $\varphi(a) = 0$. For even positive integers k, $\varphi(a) = C(a_6)^k$ for some constant $C = C_k$, where $a = \sum_{i=1}^{12} a_iY_i$.*

This gives the following corollary, which can also be checked directly.

Corollary 6.11. *For any even positive integer k and any constant C*

$$\varphi(a) = C(a_6)^k$$

is an $\mathrm{Ad}_{\tilde{G}}$-invariant polynomial on δ.

Proof. Let $\pi_6 : \delta \to \mathbb{C}$ be the projection given by

$$\pi_6\left(\sum_{j=1}^{12} a_iY_i\right) = a_6. \tag{6.1}$$

Since

$$\pi_6(\mathrm{Ad}_{\tilde{G}}(\sum_{j=1}^{12} a_iY_i)) = a_6,$$

we see that $\varphi(a) = C(a_6)^k$ is an Ad-invariant polynomial on δ. □

Example 1. In the notation of (3.2), for integers M and N we get Casimir elements on $L\delta$

$$\varphi_{m,n}\left(\sum_{i=M}^{N}\sum_{j=1}^{12} L_i^jY_j\lambda^i\right) = \mathrm{Res}_{\lambda=0}(\lambda^{-n}C(\lambda^m L_6^i(\lambda^i))^{2k})$$

$$= C\sum_S \begin{pmatrix} & & k & \\ i_M & i_{M+1} & \cdots & i_N \end{pmatrix} (L_M^6)^{i_M}\cdots(L_N^6)^{i_N}$$

for all nonnegative integers m, n, where

$$S = \{(i_M, \ldots, i_N) \mid i_M \geq 0, \ldots, i_N \geq 0,\ i_M + i_{M+1} + \ldots + i_N = k,$$

$$Mi_M + (M+1)i_{M+1} + \ldots + Ni_N = -1 + n - km\}.$$

Example 2. The natural pairing $\langle\cdot,\cdot\rangle$ on δ induces an Ad-invariant polynomial

$$\psi(Y) = \langle Y, Y\rangle = 2\sum_{i=1}^{6} a_i a_{i+6}$$

for $Y = \sum_{i=1}^{12} a_i Y_i$. Then $\psi_{m,n} : L\delta \to \mathbb{C}$ given by (3.2) becomes

$$\psi_{m,n}(L(\lambda)) = 2\sum_{j=1}^{6} \sum_{i+k-n+2m=-1} L_i^j L_k^{j+6}.$$

Example 3. (3.4) gives other Ad-invariant polynomials on $L\delta$. An explicit computation gives

$$\psi^k_{m,n}(L(\lambda)) = \sum_S \begin{pmatrix} k \\ \{i_{a,b}\}_{a,b\in\{-M,\dots,N\}} \end{pmatrix} \prod_{a,b\in\{-M,\dots,N\}} (2\sum_{j=1}^{6} L_a^j L_b^{j+6})^{i_{a,b}},$$

where

$$S = \{\{i_{a,b}\}_{a,b\in\{-M,\dots,N\}} \mid \sum_{a,b\in\{-M,\dots,N\}} i_{a,b} = k \ \text{ and } \sum_{a,b\in\{-M,\dots,N\}} i_{a,b}(a+b) - n = -1\}.$$

6.5. **The Lax pair equation in local coordinates.** We can write the Lax pair equation

$$\frac{dL}{dt} = [M, L], \quad M = \frac{1}{2}R(I(d\kappa(L(\lambda))) \tag{6.2}$$

in local coordinates when the Casimir function κ is $\varphi_{m,n}$ or $\psi_{m,n}$ given above.

Note that

$$M = \frac{1}{2}R(I(d\kappa(L(\lambda))) = \frac{1}{2}\frac{\partial(\kappa(L(\lambda))}{\partial L_p^t} Y_t \lambda^{-1-p} r(-1-p), \tag{6.3}$$

where $r(s) = 1$ if $s \geq 0$ and $r(s) = -1$ if $s < 0$.

The case $\kappa = \varphi_{m,n}$.

Lemma 6.12. *In local coordinates the Lax pair equation (6.2) becomes*

$$\frac{dL^j_{i+p}}{dt} = 0.$$

for $0 \leq j \leq 12$. Thus $L(t) = L(0)$ for any t.

Proof.

$$\frac{\partial\varphi_{mn}(\sum_{i=M}^{N}\sum_{j=1}^{12} L_i^j Y_j \lambda^i)}{\partial L_p^6} = C\sum_S \begin{pmatrix} & & k & \\ i_M & i_{M+1} & \cdots & i_N \end{pmatrix} (L_M^6)^{i_M} \cdots ((i_p)(L_p^6)^{i_p-1}) \cdot (L_N^6)^{i_N},$$

where S is some set of multi-indices. Then

$$M = R(I(\frac{\partial\varphi_{mn}(\sum_{i=M}^{N}\sum_{j=1}^{12} L_i^j Y_j \lambda^i)}{\partial L_p^6})Y_6\lambda^p) = r(-1-p)f(L(\lambda))Y_{12}.$$

Since $\mathrm{ad}(Y_{12}) = 0$, we have $[L, M] = 0$, and so $\frac{dL(t)}{dt} = 0$. $\square$

Thus the Lax pair equation is trivial in this case.

The case $\kappa = \psi_{m,n}$. By Proposition 4.3, $M = R(X) = R(\lambda^{-n+2m}L(\lambda))$.

Let q_{ij}^k be the structure constants of δ in the usual basis.

Theorem 6.13. *In local coordinates the Lax pair equation (6.2) becomes*

$$\frac{dL^k_{i+p-n+2m}}{dt} = r(-n+2m+p)\sum_{j=1}^{12}\sum_{t=1}^{12} L_i^j L_p^t q_{tj}^k$$

for $j \in \{1, \ldots, 12\}$ *and all* i, p, *where* $r(s) = 1$ *if* $s \geq 0$ *and* $r(s) = -1$ *if* $s < 0$.

Note that the Lax pair equation $\pi_1(\dot{L}(t)) = [\pi_1(L), \pi_1(M)]$ on $L\mathfrak{g}_0$ given by (4.3) has the local coordinate form

$$\frac{dL^k_{i+p-n+2m}}{dt} = r(-n+2m+p)\sum_{j=1}^{6}\sum_{t=1}^{6} L_i^j L_p^t c_{tj}^k$$

for $j \in \{1, \ldots, 6\}$ and all i, p.

6.6. **The Birkhoff factorization of** $\exp(-tX)$**.** We compute the factorization of $\exp(-tX)$ for the interesting case of $X = I(d\psi_{m,n}(L(\lambda)))$, for $\pi_6(L(\lambda)) = 0$, where π_6 is the extension to $L\delta$ of π_6 in (6.1). Then $X = \lambda^{-n+2m}L(\lambda) = \sum_{j,i}(L_i^j\lambda^{i-n+2m})Y_j$, and

$$\exp(-tX) = \exp(\sum_{j=1}^{12}(-\sum_i L_i^j\lambda^{i-n+2m}t)Y_j).$$

Let $z_j = -\sum_i L_i^j\lambda^{i-n+2m}$ for $j \in \{1, \ldots, 12\}$.

Our assumption is that $z_6 = 0$, as the exponential of $L(\lambda)$ has a simpler form in this case; in fact this is the only case needed for our main theorem below. The exponential of $L\delta$ on $z_6 = 0$ is given by

$$\exp(tz_1, tz_2, tz_3, tz_4, tz_5, 0, tz_7, tz_8, tz_9, tz_{10}, tz_{11}, tz_{12}) =$$

$$\Big(tz_1, tz_2, tz_3, tz_4, tz_5, 0, t^2z_{10}z_2 + tz_7, -t^2z_1z_{10} + tz_8, tz_9, tz_{10}, tz_{11},$$
$$tz_{12} - \frac{1}{3}t^3z_1z_{10}z_2 + \frac{3}{2}t^2z_{10}z_4 + 2t^2z_{11}z_5 + \frac{1}{2}t^2z_1z_7 + t^2z_2z_8 + \frac{3}{2}t^2z_3z_9\Big).$$

By Lemma 5.4, the Birkhoff decomposition $(g, \alpha) = (g_-, \alpha_-)^{-1}(g_+, \alpha)_+$, with $g \in G_0$ and $\alpha \in \mathfrak{g}_0^*$ is given by $g = g_-^{-1}g_+$, and $-\alpha_- + \alpha_+ = \mathrm{Ad}^*(g_-)\alpha$.

Let g_{j-} and g_{j+}, $j \in \{1, \ldots, 12\}$, be the components of g_- and g_+ respectively. Therefore, for $j \in \{1, 2, 3, 4, 5, 9, 10, 11\}$, we have

$$g_{j+} = -\sum_{i \geq n-2m} L_i^j\lambda^{i-n+2m}t$$

and

$$g_{j-} = -\sum_{i<n-2m} L_i^j \lambda^{i-n+2m} t. \tag{6.4}$$

Then

$$g_{i+} = P_+(tz_i), \quad g_{i-} = -P_-(tz_i) \tag{6.5}$$

for $i \in \{1, 2, 3, 5\}$. We also get

$$g_{4+} = P_+(tz_4 + t^2(P_-(z_1)P_+(z_2) - P_-(z_1)P_+(z_2))), \tag{6.6}$$

$$g_{4-} = P_-(tz_4 + t^2(P_-(z_1)P_+(z_2) - P_-(z_1)P_+(z_2))), \tag{6.7}$$

$$g_{6-} = 0, \qquad g_{6+} = 0, \tag{6.8}$$

$$g_{7-} = -P_-(2tg_{2-}z_{10} + t^2 z_{10} z_2 + tz_7), \tag{6.9}$$

$$g_{8-} = -P_-(-2tg_{1-}z_{10} - t^2 z_1 z_{10} + tz_8), \tag{6.10}$$

$$\begin{aligned} g_{12-} \;=\; & -P_-(-tg_{1-}g_{2-}z_{10} + 3tg_{4-}z_{10} - 2t^2 g_{2-}z_1 z_{10} + 4tg_{5-}z_{11} \\ & +tz_{12} + t^2 g_{1-}z_{10}z_2 - \frac{1}{3}t^3 z_1 z_{10} z_2 + \frac{3}{2}t^2 z_{10} z_4 + 2t^2 z_{11} z_5 + tg_{1-}z_7 \\ & +\frac{1}{2}t^2 z_1 z_7 + 2tg_{2-}z_8 + t^2 z_2 z_8 + 3tg_{3-}z_9 + \frac{3}{2}t^2 z_3 z_9), \end{aligned} \tag{6.11}$$

$$g_{k+} = z_k + g_{k-},$$

for $k \in \{7, 8, 12\}$. Then $g_- = (g_{1-}, \ldots, g_{12-})$ and $g_+ = (g_{1+}, \ldots, g_{12+})$ satisfy $g_-^{-1} g_+ = \exp(-tX)$.

We now assemble the final formulas needed to compute the solution to the Lax pair equation given by Theorem 4.1. Let $\pi : \delta \to \mathfrak{g}$ and $\theta : \delta \to \mathfrak{g}^*$ be the projections onto $\mathfrak{g}$ respectively $\mathfrak{g}^*$. Then

(6.12)

$$\pi(\mathrm{Ad}^*(g_{1-}, \ldots, g_{12-})L(\lambda))) = \mathrm{Ad}^*_{G_0}(g_{1-}, \ldots, g_{6-})\pi(L(\lambda))) + H(g_{1-}, \ldots, g_{12-})\theta(L(\lambda))$$

and

$$\theta(\mathrm{Ad}^*(g_{1-}, \ldots, g_{12-})L(\lambda))) = \mathrm{Ad}_{G_0}(g_{1-}, \ldots, g_{6-})\theta(L(\lambda))), \tag{6.13}$$

where H is given by Proposition 6.9. (It is fortunate that $\mathrm{Ad}^*_{\tilde{G}}$ does not depend on g_{12-}, which by (6.11) is difficult to compute explicitly.)

6.7. **The Feynman rules characters and the main result.** For any graph γ, the Feynman rules integral $\varphi(\lambda, q)(\gamma)$ can be written in term of Γ-functions. Some explicit formulas are as follows.

Lemma 6.14.

$$\begin{aligned}
\varphi(\lambda)(A) &= \pi^3(q^2)^{1-\lambda}B_1(\lambda)\\
\varphi(\lambda)(AA) &= \pi^6(q^2)^{1-2\lambda}B_1(\lambda)B_2(\lambda)\\
\varphi(\lambda)(AAA) &= \pi^9(q^2)^{1-3\lambda}B_1(\lambda)B_2(\lambda)B_3(\lambda)\\
\varphi(\lambda)(AAAA) &= \pi^{12}(q^2)^{1-4\lambda}B_1(\lambda)B_2(\lambda)B_3(\lambda)B_4(\lambda)\\
\varphi(\lambda)(B) &= \pi^9(q^2)^{1-3z}B_1(\lambda)^2B_2(\lambda)
\end{aligned}$$

where $B_j(\lambda) = \frac{-1}{j\lambda(1-j\lambda)(2-j\lambda)(3-j\lambda)}$, $j \in \{1, 2, 3, 4\}$.

Theorem 6.15. *Let* H_0 *be the Hopf subalgebra generated by* A, AA, AAA, $AAAA$, B, $\emptyset$, *and choose a character* $\bar\varphi(\lambda) \in \Omega\tilde{G}$ *with* $\pi_6(\bar\varphi) = 0$. *Set*

$$L_0(\lambda) = \frac{1}{2}\lambda^{n-2m}\exp^{-1}(\bar\varphi(\lambda)), \quad X = I(d\psi_{m,n}(L_0(\lambda)),$$

where $\psi_{m,n}$ *is the Casimir function on* $\Omega\delta$ *given by*

$$\psi_{m,n}(L(\lambda))) = \mathrm{Res}_{\lambda=0}(\lambda^m\langle\lambda^n L(\lambda), \lambda^n L(\lambda)\rangle).$$

Then $\exp(X) = \bar\varphi$ *and the solution of*

$$\frac{dL}{dt} = [M, L], \quad M = \frac{1}{2}R(I(d\psi_{m,n}(L(\lambda))) \tag{6.14}$$

with the initial condition $L(0) = L_0$ *is given by*

$$L(t) = \mathrm{Ad}^* g_\pm(t)\cdot L_0, \tag{6.15}$$

with $\exp(-tX) = g_-(t)^{-1}g_+(t)$ *where* g_- *and* g_+ *are given by (6.4), (6.8), (6.9), (6.10), (6.11), and where the* $\mathfrak{g}_0$ *and* $\mathfrak{g}_0^*$ *components of* Ad *are given by (6.12) and (6.13).*

In the particular case where $\bar\varphi$ *is the normalized Feynman rule characters* $\bar\varphi(\lambda) = \frac{\varphi(\lambda)}{q^2}$, $g_-(t)$ *and* $g_+(t)$ *are given by (6.5), (6.6) and (6.7), and the solution* $L(t)$ *of (6.14) is the flow of Feynman rules.*

This follows from Theorem 5.6.

Remark 6.16. To any Lax equation with a spectral parameter, one can associate a spectral curve and study its algebro-geometric properties (see [8]). In our case, we consider the adjoint representation ad : $\delta \to \mathfrak{gl}(\delta)$ and the induced adjoint representation of the loop algebra. The spectral curve is given by the characteristic equation of $\mathrm{ad}(L\lambda)$ $\Gamma_0 = \{(\lambda, \nu) \in \mathbb{C} - \{0\} \times \mathbb{C} \mid \det(\mathrm{ad}(L(\lambda) - \nu\mathrm{Id}) = 0\}$.

The theory of the spectral curve and its Jacobian usually assumes that the spectral curve is irreducible. Unfortunately, for all 43 bialgebra structures on δ, on the associated

Lie algebra ad(δ) all eigenvalues of the characteristic equation are zero, and the zero eigenspace is nine dimensional. The spectral curve itself is the union of degree one curves. Thus each irreducible component has a trivial Jacobian, and the spectral curve theory breaks down.

References

1. M. Adler, P. van Moerbeke, and P. Vanhaecke, *Algebraic integrability, Painlevé geometry and Lie algebras*, Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics, vol. 47, Springer-Verlag, Berlin, 2004.
2. C. Chryssomalakos, H. Quevedo, M. Rosenbaum, and J. D. Vergara, *Normal Coordinates and Primitive Elements in the Hopf Algebra of Renormalization*, Comm. Math. Phys. **225** (2002), 465–485, hep–th/0105259.
3. A. Connes and D. Kreimer, *Renormalization in Quantum Field Theory and the Riemann-Hilbert Problem I: The Hopf algebra structure of Graphs and the Main Theorem*, Comm. Math. Phys. **210** (2000), 249–273.
4. K. Ebrahimi-Fard, L. Guo, and D. Kreimer, *Integrable renormalization ii: the general case*, Annales Inst. Henri Poincaré **6** (2005), 369–395,hep–th/0403118.
5. D. Kastler, *Connes-Moscovici-Kreimer Hopf algebras*, Mathematical Physics in Mathematics and Physics (Sienna, 2000), Fields Inst. Commun., vol. 30, Amer. Math. Soc., Providence, RI, 2001, pp. 219–248, hep–th/0105259.
6. Y. Kosmann-Schwarzbach, *Lie Bialgebras, Poisson Lie Groups and Dressing Transformations*, Springer-Verlag, Berlin Heidelberg New York, 1997.
7. D. Manchon, *Hopf algebras, from basics to applications to renormalization*, Comptes-rendus des Rencontres mathématiques de Glanon 2001, 2003, math.QA/0408405.
8. A.G. Reyman and M.A. Semenov-Tian-Shansky, *Integrable Systems II: Group-Theoretical Methods in the Theory of Finite-Dimensional Integrable Systems*, Dynamical systems. VII, Encyclopaedia of Mathematical Sciences, vol. 16, Springer-Verlag, Berlin, 1994.

DEPARTMENT OF MATHEMATICS AND STATISTICS, BOSTON UNIVERSITY, BOSTON, MA 02215.
`baditoiu@math.bu.edu`

DEPARTMENT OF MATHEMATICS AND STATISTICS, BOSTON UNIVERSITY, BOSTON, MA 02215.
`sr@math.bu.edu`